



Restriction Enzyme

Alw I



Cat.# FG-AlwI	Size 500 units	Conc. 10 units/μl
-------------------------	--------------------------	-----------------------------

Store at -20°C

Supplied with: 10X FastGene® Buffer IV (FG-REB4)
10X FastGene® FastCut Buffer (FG-REBHF)
6X DNA Loading Buffer
Sterile water

Recognition site



For Research Use Only. Not for use in diagnostic procedures.



Source

Acinetobacter lwoffii

Reaction conditions

- 1X FastGene® Buffer IV, 37°C
- 1X FastGene® FastCut Buffer, 37°C

FastGene® FastCut Buffer

FastGene® restriction enzyme can cut substrate DNA in 5-15 min with FastGene® FastCut Buffer.

1X FastGene® Buffer IV

- 20 mM Tris-acetate (pH 7.9 at 25°C)
- 50 mM potassium acetate
- 10 mM magnesium acetate
- 100 μg/ml BSA

Unit definition

One unit is defined as the amount of enzyme required to digest 1 μg of Lambda DNA (dam-) in 1 hour at 37°C in a total reaction volume of 50 μl.

Quality control

- Unit definition assay
- Overdigestion assay
- Endonuclease assay
- Extreme pure assay

Dilution buffer

FastGene® Diluent A

Heat Inactivation

No.

Methylation sensitivity

- dam* methylation: Sensitive
- dcm* methylation: Not sensitive
- CpG methylation: Not sensitive

Relative activity in FastGene® Buffers

FastGene® Buffer I:	50%
FastGene® Buffer II:	50%
FastGene® Buffer III:	10%
FastGene® Buffer IV:	100%
FastGene® FastCut Buffer:	100%

Note

DNA cleavage is blocked by dam methylation. It produces a 5' extension of one nucleotide, which is more difficult to be ligated than blunt-ends. Reaction condition with excess enzyme, excess glycerol (>5%) or extended digestion may result in star activity.

Standard reaction condition

- Normal protocol

Component	Final Conc.	Volume
Substrate DNA	1 μg	X μl
10X FastGene® Buffer IV	1 X	5 μl
Alw I	Substrate dependent	
Sterile water		up to 50 μl
→ Incubate at 37°C for 1 hr		

- Fast protocol

Component	Final Conc.	Volume
Substrate DNA	1 μg	X μl
10X FastGene® FastCut Buffer	1 X	5 μl
Alw I	10 unit	1 μl
Sterile water		up to 50 μl
→ Incubate at 37°C for 15 min		

※ We recommend 5-10 units of enzyme per μg DNA and 10-20 units for genomic DNA in a 1 h digest.