



Restriction Enzyme Bal I



Cat.# FG-Bal	Size 50 units	Conc. 5 units/μl
-----------------	------------------	---------------------

Store at -20°C

Supplied with: 10X FastGene® Buffer Bal I (FG-REBBal)
6X DNA Loading Buffer
Sterile water

Recognition site



For Research Use Only. Not for use in diagnostic procedures.

ISO9001

Dilution buffer:

FastGene® Diluent B

Heat Inactivation

Bal I can be inactivated at 65°C for 20 min.

Methylation sensitivity

dam methylation: Not sensitive

dcm methylation: Conditionally sensitive

CpG methylation: Not sensitive

Relative activity in FastGene® Buffers

FastGene® Buffer I:	0%
FastGene® Buffer II:	75%
FastGene® Buffer III:	25%
FastGene® Buffer IV:	75%

Note

Activity is inhibited by *dcm* methylation partially overlapping its recognition sequence.

Source: *Brevibacterium albidum*

Reaction conditions

1X FastGene® Buffer Bal I 37°C

1X FastGene® Buffer Bal I

50 mM Tris-HCl (pH 8.2 at 25°C)

5 mM MgCl₂

Unit definition

One unit is defined as the amount of enzyme required for complete digestion of 1 μg bacteriophage λ at 37°C for 1 hr in 50 μl reaction mixtures.

Quality control

- Unit definition assay
- Overdigestion assay
- Endonuclease assay
- Extreme pure assay

Standard reaction condition

- Normal protocol

Component	Final Conc.	Volume
Substrate DNA	1 μg	X μl
10X FastGene® Buffer Bal I	1 X	5 μl
Bal I	5 unit	1 μl
Sterile water		up to 50 μl

→ Incubate at 37°C for 1 hr

※ We recommend 5-10 units of enzyme per μg DNA and 10-20 units for genomic DNA in a 1 h digest.