



## Restriction Enzyme PspG I




Cat.#	Size	Conc.
FG-PspGI	1,000 units	10 units/μl

Store at -20°C

**Supplied with:** 10X FastGene® Buffer IV (FG-REB4)  
6X DNA Loading Buffer  
Sterile water

### Recognition site



For Research Use Only. Not for use in diagnostic procedures.

ISO9001

**Source:** *Pyrococcus* species strain GI-H

### Reaction conditions

1X FastGene® Buffer IV 37°C or 75°C

### 1X FastGene® Buffer IV

20 mM Tris-acetate (pH 7.9 at 25°C)  
50 mM potassium acetate  
10 mM magnesium acetate  
100 μg/ml BSA

### Unit definition

One unit is defined as the amount of enzyme required to digest 1 μg of T7 DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

### Quality control

- Unit definition assay
- Overdigestion assay
- Endonuclease assay
- Extreme pure assay

### Dilution buffer:

FastGene® Diluent A

### Heat Inactivation

No.

### Methylation sensitivity

*dam* methylation: Not sensitive  
*dcm* methylation: sensitive  
CpG methylation: Not sensitive

### Prolonged incubation

A minimum amount of enzyme required to digest 1 μg substrate DNA for 16 hr; 0.13 U.

### Relative activity in FastGene® Buffers

FastGene® Buffer I: 25%  
FastGene® Buffer II: 100%  
FastGene® Buffer III: 75%  
FastGene® Buffer IV: 100%  
FastGene® FastCut Buffer: Not recommended

### Standard reaction condition

- Normal protocol

Component	Final Conc.	Volume
Substrate DNA	1 μg	X μl
10X FastGene® Buffer IV	1 X	5 μl
PspG I	10 unit	1 μl
Sterile water		up to 50 μl

→ Incubate at 37°C or 75°C for 1 hr

※ We recommend 5-10 units of enzyme per μg DNA and 10-20 units for genomic DNA in a 1 h digest.



## Restriction Enzyme PspG I




Cat.#	Size	Conc.
FG-PspGI	1,000 units	10 units/μl

Store at -20°C

**Supplied with:** 10X FastGene® Buffer IV (FG-REB4)  
6X DNA Loading Buffer  
Sterile water

### Recognition site



For Research Use Only. Not for use in diagnostic procedures.

ISO9001

**Source:** *Pyrococcus* species strain GI-H

### Reaction conditions

1X FastGene® Buffer IV 37°C or 75°C

### 1X FastGene® Buffer IV

20 mM Tris-acetate (pH 7.9 at 25°C)  
50 mM potassium acetate  
10 mM magnesium acetate  
100 μg/ml BSA

### Unit definition

One unit is defined as the amount of enzyme required to digest 1 μg of T7 DNA in 1 hour at 37°C in a total reaction volume of 50 μl.

### Quality control

- Unit definition assay
- Overdigestion assay
- Endonuclease assay
- Extreme pure assay

### Dilution buffer:

FastGene® Diluent A

### Heat Inactivation

No.

### Methylation sensitivity

*dam* methylation: Not sensitive  
*dcm* methylation: sensitive  
CpG methylation: Not sensitive

### Prolonged incubation

A minimum amount of enzyme required to digest 1 μg substrate DNA for 16 hr; 0.13 U.

### Relative activity in FastGene® Buffers

FastGene® Buffer I: 25%  
FastGene® Buffer II: 100%  
FastGene® Buffer III: 75%  
FastGene® Buffer IV: 100%  
FastGene® FastCut Buffer: Not recommended

### Standard reaction condition

- Normal protocol

Component	Final Conc.	Volume
Substrate DNA	1 μg	X μl
10X FastGene® Buffer IV	1 X	5 μl
PspG I	10 unit	1 μl
Sterile water		up to 50 μl

→ Incubate at 37°C or 75°C for 1 hr

※ We recommend 5-10 units of enzyme per μg DNA and 10-20 units for genomic DNA in a 1 h digest.