



Restriction Enzyme Pvu I



| Cat.# | Size | Conc. |
|---------|-----------|-------------|
| FG-PvuI | 500 units | 10 units/μl |

Store at -20°C

Supplied with: 10X FastGene® Buffer III (FG-REB3)
10X FastGene® FastCut Buffer (FG-REBHF)
6X DNA Loading Buffer
Sterile water

Recognition site



For Research Use Only. Not for use in diagnostic procedures.

ISO9001

Source: *Proteus vulgaris*

Reaction conditions

1X FastGene® Buffer III 37°C
1X FastGene® FastCut Buffer, 37°C

FastGene® FastCut Buffer

FastGene® restriction enzyme can cut substrate DNA in 5-15 with FastGene® FastCut Buffer.

1X FastGene® Buffer III

50 mM Tris-HCl (pH 7.9 at 25°C)
100 mM NaCl
10 mM MgCl₂
100 μg/ml BSA

Unit definition

One unit is defined as the amount of enzyme required for complete digestion of 1 μg bacteriophage λ at 37°C for 1 hr in 50 μl reaction mixtures.

Quality control

- Unit definition assay
- Overdigestion assay
- Endonuclease assay
- Extreme pure assay

Dilution buffer:

FastGene® Diluent B

Heat Inactivation

No.

Methylation sensitivity

dam methylation: Not sensitive
dcm methylation: Not sensitive
CpG methylation: sensitive

Prolonged incubation

A minimum amount of enzyme required to digest 1 μg substrate DNA for 16 hr; 0.13 U.

Relative activity in FastGene® Buffers

| | |
|---------------------------|------|
| FastGene® Buffer I: | 25% |
| FastGene® Buffer II: | 75% |
| FastGene® Buffer III: | 100% |
| FastGene® Buffer IV: | 50% |
| FastGene® FastCut Buffer: | 100% |

Note

Cleavage of mammalian genomic DNA is blocked by CpG methylation.

Standard reaction condition

- Normal protocol

| Component | Final Conc. | Volume |
|--------------------------|-------------|-------------|
| Substrate DNA | 1 μg | X μl |
| 10X FastGene® Buffer III | 1 X | 5 μl |
| Pvu I | 10 unit | 1 μl |
| Sterile water | | up to 50 μl |

→ Incubate at 37°C for 1 hr

- Fast protocol

| Component | Final Conc. | Volume |
|------------------------------|-------------|-------------|
| Substrate DNA | 1 μg | X μl |
| 10X FastGene® FastCut Buffer | 1 X | 5 μl |
| Pvu I | 10 unit | 1 μl |
| Sterile water | | up to 50 μl |

→ Incubate at 37°C for 15 min

※ We recommend 5-10 units of enzyme per μg DNA and 10-20 units for genomic DNA in a 1 h digest.



Restriction Enzyme Pvu I



| Cat.# | Size | Conc. |
|---------|-----------|-------------|
| FG-PvuI | 500 units | 10 units/μl |

Store at -20°C

Supplied with: 10X FastGene® Buffer III (FG-REB3)
10X FastGene® FastCut Buffer (FG-REBHF)
6X DNA Loading Buffer
Sterile water

Recognition site



For Research Use Only. Not for use in diagnostic procedures.

ISO9001

Source: *Proteus vulgaris*

Reaction conditions

1X FastGene® Buffer III 37°C
1X FastGene® FastCut Buffer, 37°C

FastGene® FastCut Buffer

FastGene® restriction enzyme can cut substrate DNA in 5-15 with FastGene® FastCut Buffer.

1X FastGene® Buffer III

50 mM Tris-HCl (pH 7.9 at 25°C)
100 mM NaCl
10 mM MgCl₂
100 μg/ml BSA

Unit definition

One unit is defined as the amount of enzyme required for complete digestion of 1 μg bacteriophage λ at 37°C for 1 hr in 50 μl reaction mixtures.

Quality control

- Unit definition assay
- Overdigestion assay
- Endonuclease assay
- Extreme pure assay

Dilution buffer:

FastGene® Diluent B

Heat Inactivation

No.

Methylation sensitivity

dam methylation: Not sensitive
dcm methylation: Not sensitive
CpG methylation: sensitive

Prolonged incubation

A minimum amount of enzyme required to digest 1 μg substrate DNA for 16 hr; 0.13 U.

Relative activity in FastGene® Buffers

| | |
|---------------------------|------|
| FastGene® Buffer I: | 25% |
| FastGene® Buffer II: | 75% |
| FastGene® Buffer III: | 100% |
| FastGene® Buffer IV: | 50% |
| FastGene® FastCut Buffer: | 100% |

Note

Cleavage of mammalian genomic DNA is blocked by CpG methylation.

Standard reaction condition

- Normal protocol

| Component | Final Conc. | Volume |
|--------------------------|-------------|-------------|
| Substrate DNA | 1 μg | X μl |
| 10X FastGene® Buffer III | 1 X | 5 μl |
| Pvu I | 10 unit | 1 μl |
| Sterile water | | up to 50 μl |

→ Incubate at 37°C for 1 hr

- Fast protocol

| Component | Final Conc. | Volume |
|------------------------------|-------------|-------------|
| Substrate DNA | 1 μg | X μl |
| 10X FastGene® FastCut Buffer | 1 X | 5 μl |
| Pvu I | 10 unit | 1 μl |
| Sterile water | | up to 50 μl |

→ Incubate at 37°C for 15 min

※ We recommend 5-10 units of enzyme per μg DNA and 10-20 units for genomic DNA in a 1 h digest.