FastGene®

Restriction Enzyme
Fsp I

IV · 37℃ · 65 · CpG

Cat.#	Size	Conc.
FG-FspI	500 units	5 units/µl

Store at -20℃

Supplied with: 10X FastGene® Buffer IV (FG-REB4)
10X FastGene® FastCut Buffer (FG-REBHF)
6X DNA Loading Buffer
Sterile water

Recognition site

```
5' ··· T G C│G C A ··· 3'
3' ··· A C G│C G T ··· 5'
```

For Research Use Only. Not for use in diagnostic procedures.

ISO9001

Source: *Fischerella* species

Reaction conditions
1X FastGene® Buffer IV 37℃
1X FastGene® FastCut Buffer, 37℃

FastGene® FastCut Buffer
FastGene® restriction enzyme can cut substrate DNA in 5-15 with FastGene® FastCut Buffer.

1X FastGene® Buffer IV
20 mM Tris-acetate (pH 7.9 at 25℃)
50 mM potassium acetate
10 mM magnesium acetate
100 µg/ml BSA

Unit definition
One unit is defined as the amount of enzyme required for complete digestion of 1 µg bacteriophage λ at 37℃ for 1 hr in 50 µl reaction mixtures.

Quality control
- Unit definition assay
- Overdigestion assay
- Endonuclease assay
- Extreme pure assay

Dilution buffer:
FastGene® Diluent C

Heat Inactivation
No heat inactivation.

Methylation sensitivity
dam methylation: Not sensitive
dcm methylation: Not sensitive
CpG methylation: sensitive

Prolonged incubation
A minimum amount of enzyme required to digest 1 µg substrate DNA for 16 hr; 0.13 U.

Relative activity in FastGene® Buffers
FastGene® Buffer I:	75%
FastGene® Buffer II:	100%
FastGene® Buffer III:	50%
FastGene® Buffer IV:	100%
FastGene® FastCut Buffer:	100%

Note
Cleavage of mammalian genomic DNA is blocked by CpG methylation.

Standard reaction condition
- Normal protocol

Component	Final Conc.	Volume
Substrate DNA	1 µg	X µl
10X FastGene® Buffer IV	1 X	5 µl
Fsp I	5 unit	1 µl
Sterile water		up to 50 µl

→ Incubate at 37℃ for 1 hr

- Fast protocol

Component	Final Conc.	Volume
Substrate DNA	1 µg	X µl
10X FastGene® FastCut Buffer	1 X	5 µl
Fsp I	5 unit	1 µl
Sterile water		up to 50 µl

→ Incubate at 37℃ for 15 min

※ We recommend 5-10 units of enzyme per µg DNA and 10-20 units for genomic DNA in a 1 h digest.